

I wish to express my thanks and appreciation to Dr. Harlow B. Mills, for his help and suggestions, and to Dr. J. Bequaert, Curator of Insects, MCZ, Harvard University, Cambridge, Mass., for his kindness and permission to work there examining the types of various species.

LITERATURE USED AND CITED

- Banks, Nathan.** 1903. New Smythuridae from the District of Columbia. Proceedings of the Entomological Society of Washington, 5: 154-155. (Describes *Dicyrtoma frontalis* Banks).
- Folsom, J. W.** 1896a. New Species of Papirius. Psyche 7: 344-345. (Describes *Dicyrtoma hageni* Folsom originally under *Papirius hagenii* Folsom).
- . 1934. Redescriptions of North American Sminthuridae. Iowa State College Journal of Science, vol. VIII, no. 4, pp. 461-511. (Redescribes *Papirius hagenii* Folsom under *Dicyrtoma hageni* Folsom).
- Mills, Harlow B.** 1934. A Monograph of the Collembola of Iowa. Collegiate Press, Inc., Ames, Iowa, 143 pp., 12 Pls. (Describes *Dicyrtoma quadangularis* Mills).

Rhagionidae (Diptera) from the Great Smoky Mountains National Park. Five species of this family were collected by the writer and Robert R. Dreisbach in the Tennessee part of the park in 1946: *Chrysopilus connexus* Jns., Elkmont, June 15, 2 ♀♀; *C. ornatus* Say, Cherokee Orchard, June 14, ♀; *C. quadratus* Say, Elkmont, June 15, ♀; *C. rotundipennis* Lw., Cades Cove, June 13, ♂; *Rhagio vertebratus* Lw., Cherokee Orchard, June 11, Chimneys Camp, June 11, Headquarters, June 15, 3 ♂♂, 3 ♀♀. It is interesting to note that the only form found in Cades Cove (*Chrysopilus rotundipennis*) has a costal plains distribution. *Chrysopilus connexus* is apparently known only from the North Carolina mountains and Jacksonville, Florida. The others are more widespread.

—GEORGE C. STEYSKAL, Grosse Ile, Michigan.